

# Tuesday 20 October 2020 - Afternoon

## A Level Biology A

H420/03 Unified biology

Time allowed: 1 hour 30 minutes

# \* 8 2 4 3 2 1 2 7 0 9

You	can	use:
-----	-----	------

- · a scientific or graphical calculator
- a ruler (cm/mm)



Please write clearly in black ink. <b>Do not write in the barcodes.</b>								
Centre number					Candidate number			
First name(s)								
Last name								

#### **INSTRUCTIONS**

- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided. If you need extra space use the lined pages at the end of this booklet. The question numbers must be clearly shown.
- · Answer all the questions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.

#### **INFORMATION**

- The total mark for this paper is 70.
- The marks for each question are shown in brackets [ ].
- Quality of extended response will be assessed in questions marked with an asterisk (\*).
- This document has 28 pages.

#### **ADVICE**

· Read each question carefully before you start your answer.

## Answer **all** the questions.

1 (a) A student dissected a kidney. Fig. 1.1 shows one half of the dissected kidney.



Fig. 1.1

Draw a simple diagram of the kidney in Fig. 1.1 in the space below.

On your diagram, label the pelvis, medulla and cortex.

(b) A photomicrograph of a stained section of kidney tissue is shown in Fig. 1.2.




Fig. 1.2

(i)	State one function of the distal convoluted tubule.
	[1]
(ii)	Describe the function of the structures labelled <b>A</b> in Fig. 1.2.
	[1]
(iii)	Suggest which lumen, ${\bf B}$ or ${\bf C}$ , has the highest concentration of urea. Explain your answer.
	[1]

 $(c)^*$  Water reabsorption in the kidney is controlled by the endocrine and nervous systems.

Aldosterone and	ADH are horm	ones	that act o	n the kidne	ey.				
Aldosterone caus	ses sodium ior	s to b	e pumped	I from the	collect	ing duct c	ells	into tissue	e fluid.
Describe how the reabsorption from				systems	work	together	to	increase	water
							• • • • • •		
							• • • • • •		
									[6]
Additional answe	r space if requ	iired.							
							• • • • • •		

(d) Diuretics are drugs that decrease the reabsorption of water into the blood from the kidney.

Diuretics can change the concentration of ions and other molecules in the blood.

Some diuretics are used to treat high blood pressure.

The table below lists three different diuretics, **X**, **Y** and **Z**, and some of their effects in the body.

	Without a			
	diuretic	Х	Υ	Z
Rate of urine production (ml min <sup>-1</sup> )	1	3	13	8
Blood chloride ion concentration (mmol dm <sup>-3</sup> )	60	15	150	150
Blood potassium ion concentration (mmol dm <sup>-3</sup> )	15	60	12	25
Blood glucose concentration (mmol dm <sup>-3</sup> )	6	6	9	8

(i)	Suggest which of the diuretics, <b>X</b> , <b>Y</b> or <b>Z</b> , would be the most effective at reducing a person's blood pressure. Explain your choice.
	diuretic
	explanation
	[1]
(ii)	Suggest which of the diuretics, <b>X</b> , <b>Y</b> or <b>Z</b> , would be the most appropriate for use by a person with type II diabetes. Explain your choice.
	diuretic
	explanation
	[1]

2 (a) Valves control the flow of blood through the heart.


Complete the table below to show the roles of two valves in the heart.

Source of blood	Valve that controls blood flow	Destination of blood
	right semilunar valve	
left atrium		left ventricle

[2]

**(b)** A ventricular septal defect (VSD) is a hole in the septum of the heart.

The diagram below shows a heart with VSD.



	Describe and explain why people with VSD can easily become tired.
	[4]
	[4]
(c)	Creatine kinase (CK) is an enzyme that catalyses reactions in heart muscle. High levels of CK in the blood indicate that a person may have had a heart attack.
	Suggest why high levels of CK in the blood indicate that a person may have had a heart attack.
	[1]
	[']
(d)	Mice are often used in laboratory studies to research treatments for heart conditions.
	These mice are often clones.
	Suggest one reason why clones are used in these studies.
	F41
	[1]

- 3 Students investigated the effect of light on the growth of garden cress seedlings.
  - A total of 120 seedlings were divided into 2 groups of 60.
  - Group A was grown in darkness for 2 days.
  - Group B was grown for 1 day in darkness and then for 1 day in white light using the set-up shown in Fig. 3.1.




Fig. 3.1

The results of the students' experiment are shown in Tables 3.1 and 3.2.

Group	Mean len	gth (mm)	Mean mass (μg)		
Group	stem	root	stem	root	
А	13	18	102	60	
В	25	23	160	120	

Table 3.1

	Number of	seedlings
Direction of growth in Group B	stem	root
Away from light	2	29
Neither away from nor towards light	3	20
Towards light	55	11

Table 3.2

(a) (i)*	Describe and explain the results shown in Tables 3.1 and 3.2.
	[6]
	Additional answer space if required.

(ii)	The students wanted to test whether there was a significant difference between the stem
	lengths of the seedlings in Group A and the seedlings in Group B.

State the name of the most appropriate statistical test for the students to use.	

(iii)	Justify your choice of statistical test given in part (ii).

[1]

.....

[1]

- (iv) Table 3.2 records the direction of growth as:
  - away from light
  - neither away from nor towards light
  - towards light.

The students used the chi-squared test to determine whether the direction of root growth was significantly different from their expectations.

Their null hypothesis was:

There is no difference between the expected direction of root growth and the observed direction of root growth.

The calculated chi-squared value was 8.10.

The students compared their chi-squared value of 8.10 to the values in Table 3.3.

Degrees of		Probability (p)					
freedom	0.10	0.05	0.01				
1	2.71	3.84	6.64				
2	4.60	5.99	9.21				
3	6.25	7.82	11.34				
4	7.78	9.49	13.28				
5	9.24	11.07	15.09				

Table 3.3

		What can the students conclude about the 8.10?	ir results based on a chi-squared value of
			[3]
(b)	The	e solution in which the seedlings were grown c	ontained various dissolved ions.
	Cor	molete the table below with the <b>formula</b> of eac	ch ion
	Cor	mplete the table below with the <b>formula</b> of each	ch ion.  Formula of ion
	Cor		
	Cor	Function Supplies elements that form part of the	
	Cor	Function  Supplies elements that form part of the structure of amino acids and nucleotides  Forms part of the structure of DNA and	
(c)		Function  Supplies elements that form part of the structure of amino acids and nucleotides  Forms part of the structure of DNA and	Formula of ion  [2]
(c)		Function  Supplies elements that form part of the structure of amino acids and nucleotides  Forms part of the structure of DNA and phospholipids	Formula of ion  [2] hormone concentration on root growth.
(c)	The	Function  Supplies elements that form part of the structure of amino acids and nucleotides  Forms part of the structure of DNA and phospholipids  e students also investigated the effect of plant  State the name of a plant hormone that woul	Formula of ion  [2] hormone concentration on root growth.
(c)	The	Function  Supplies elements that form part of the structure of amino acids and nucleotides  Forms part of the structure of DNA and phospholipids  e students also investigated the effect of plant  State the name of a plant hormone that woul	Formula of ion  [2] hormone concentration on root growth.  d be expected to affect root growth.  [1]
(c)	The	Function  Supplies elements that form part of the structure of amino acids and nucleotides  Forms part of the structure of DNA and phospholipids  e students also investigated the effect of plant  State the name of a plant hormone that woul	Formula of ion  [2] hormone concentration on root growth.  d be expected to affect root growth.  [1] ght, temperature and mineral concentration.

(d) The growth of plant roots is thought to be controlled by specialised cells called statocytes.

One hypothesis for how a statocyte controls root growth involves small organelles called amyloplasts and is shown in Fig. 3.2.

Position of root	Activity in statocyte	Effect on growth			
	falling amyloplasts endoplasmic reticulum Ca <sup>2+</sup>	direction of growth			
gravity	Ca <sup>2+</sup>	direction of growth			

Fig. 3.2

What can you conclude from the information in Fig. 3.2 about how a statocyte controls roof growth?
[2]

## 13 BLANK PAGE

PLEASE DO NOT WRITE ON THIS PAGE

**4 (a)** Prokaryotic cells have cytoskeletons. The molecules in prokaryotic cytoskeletons are different from the molecules in eukaryotic cytoskeletons.

Table 4.1 lists three molecules present in a prokaryotic cytoskeleton.




Table 4.1

(i)	Suggest the function of Protein A.
	[1]
(ii)	Suggest the function of Protein C.
	[1]

(i	(iii)	An antibiotic called A22 binds irreversibly to Protein B. Despite its antibiotic properties, A22 is not used in humans.
		Suggest why scientists have advised that A22 should not be used in humans.
		[1]
(b)		comycin is an antibiotic that has been used to treat bacterial infections for many decades reral strains of bacteria have evolved resistance to vancomycin.
	Orit	avancin is an antibiotic with a similar structure to vancomycin.

Table 4.2 shows data obtained from treatments with the two antibiotics.


oritavancin vancomycin 4 60 Years of use as an antibiotic 9.9 10.5 nausea headache 7.1 6.7 Percentage of 3.7 3.2 diarrhoea patients developing 4.6 4.7 side effects vomiting constipation 3.4 3.9 dizziness 2.7 2.6 82.5 83.5 Staphylococcus aureus Percentage of bacterial infections **MRSA** 81.4 80.6 cured Streptococcus sp. 77.2 85.3

Table 4.2

Use the data in Table 4.2 to evaluate the advantages <b>and</b> disadvantages of using oritavan rather than vancomycin as an antibiotic.	cin
	[2]

(c) Bacteria such as *E. coli* can be genetically engineered for use in medical science.

An example of the genetic engineering of *E. coli* is shown in the diagram below.




*E. coli* that have taken up the recombinant plasmid are injected into mice with liver tumours.

(i)	Complete the d	•	e by	writing	the	missing	words	or	phrases	in	the	boxes
			Ansv	ver <b>on</b> th	ne di	agram						[3]

(ii)	Suggest why the scientists used a plasmid that contained an antibiotic resistance gene.
	[1]
(iii)	The scientists observed the following:
	<ul> <li>1 in 400 bacteria took up the plasmid</li> <li>1 in 1000 of the plasmids taken up by bacteria contained the β-galactosidase gene.</li> </ul>
	Calculate the percentage of bacteria that contained the $\beta$ -galactosidase gene.
	percentage of bacteria = % [2]
(iv)	A technique called quantitative PCR is used to check that the <i>E. coli</i> population is growing on the mice liver tumours rather than on healthy tissue.
	Suggest how the scientists could use PCR to <b>compare</b> <i>E. coli</i> growth rates on cancerous liver tissue and healthy tissue.
	[2]
(v)	Some people think that the genetic engineering of certain organisms is unethical.
(*)	
	However, there are very few ethical concerns about the genetic engineering of bacteria such as <i>E. coli</i> .
	Suggest why there are very few ethical concerns about the genetic engineering of <i>E. coli</i> .
	[1]

**5** Ecologists were studying an area that contained three different habitats. The area is shown in the diagram below.



(a) The ecologists sampled the area to estimate insect biodiversity.

Describe how the ecologists should choose the number and locations of their samples to ensure that the sampling is representative.

Use a calculation to support your answer.	
	31

**(b)** Two of the insect species that were sampled were the large heath butterfly and the bog hoverfly.

The ecologists used the capture-mark-recapture technique and estimated population sizes using two different calculations: the Lincoln estimate and the Chapman estimate.

(i) Calculate the population sizes of the two insect species using each of the formulae below.

Write your answers in the table.

**Lincoln estimate formula**: population size =  $\frac{n1 \times n2}{m}$ 

Chapman estimate formula: population size =  $\left(\frac{(n1+1)\times(n2+1)}{(m+1)}\right)-1$ 

n = number of individuals in a particular sample

m = number of marked individuals in the second sample

Consider	Number captured and marked in sample 1	Total number in sample 2	Number of marked individuals in sample 2	Population estimate (number of individuals)	
Species				Lincoln estimate	Chapman estimate
large heath butterfly	77	73	4		
bog hoverfly	5	6	1		

[2]

(ii) The Lincoln and Chapman formulae give different estimates for population size.

Give **two** further conclusions about the difference in population estimates given by the Lincoln and Chapman formulae.

1	 	 	
2			

[2]

- (c) The peat bog habitat had been damaged by peat extraction and by management of the neighbouring farmland. Ecologists decided to treat the peat bog in the following way:
  - A buffer region was created between the peat bog and the neighbouring farmland.
  - No visitors were allowed on the land.
  - Ditches were blocked to raise water levels.
  - Peat extraction, tree planting and the use of fertilisers were banned.

A student suggested that this was an example of preservation.

Evaluate the student's conclusion.
[2]

(d) Conservation agreements can be national (within a particular country) or international.

Three conservation agreements are listed in the table below.

Place ticks  $(\checkmark)$  in the correct boxes to indicate which features are true for each of the three conservation agreements.

Name of agreement	International agreement	Farmers are offered payments for conservation
Environmental (Countryside) Stewardship Scheme		
Convention on International Trade in Endangered Species		
Rio Convention on Biological Diversity		

[2]

## 21 BLANK PAGE

PLEASE DO NOT WRITE ON THIS PAGE

**6 (a)** The oxygen dissociation curves for adult haemoglobin and fetal haemoglobin are shown in Fig. 6.1.




Fig. 6.1

)	haemoglobin.	auit
		. [2]

(ii) Myoglobin is a protein found in muscles. Oxygen binds to myoglobin.

A student described the oxygen dissociation curve for myoglobin as follows:

- When oxygen first becomes available, myoglobin saturation increases at a constant rate of 8% per mmHg of oxygen.
- When there is a slightly higher partial pressure of oxygen, the rate of oxygen binding slows gradually until the myoglobin is 100% saturated.
- The partial pressure at which myoglobin reaches 100% saturation is the partial pressure at which adult haemoglobin is 80% saturated.

Sketch an oxygen dissociation curve for myoglobin **on Fig. 6.1** based on the description provided above.

**(b)** Haemocyanin is an oxygen-binding pigment that is found in many invertebrate animals, including lobsters.

Fig. 6.2 shows the oxygen dissociation curves for haemoglobin and haemocyanin.




Fig. 6.2

(c)	When old red blood cells are broken down, each haem group is converted to a molecule called bilirubin. Bilirubin passes through the digestive system. Bilirubin gives faeces their characteristic colour.
	Explain why bilirubin production and processing is an example of excretion.

### **END OF QUESTION PAPER**

### **ADDITIONAL ANSWER SPACE**

If additional space is required, you should use the following lined page(s). The question number(s) must be clearly shown in the margin(s).					
	<u></u>				



	-
	• •
 	• •
	• •
	••
	• •
 	• •
 	• •
	٠.
	••
	• •
 	• •



#### Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.