

---

**PHYSICS**

**9702/21**

Paper 2 AS Level Structured Questions

**May/June 2016**

MARK SCHEME

Maximum Mark: 60

---

**Published**

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2016 series for most Cambridge IGCSE<sup>®</sup>, Cambridge International A and AS Level components and some Cambridge O Level components.

Page 2	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2016	9702	21

- 1 (a) (i)  $(50 \text{ to } 200) \times 10^{-3} \text{ kg}$  or  $(0.05 \text{ to } 0.2) \text{ kg}$  B1 [1]
- (ii)  $(50 \text{ to } 300) \text{ cm}^3$  B1 [1]
- (b) density = mass/volume or  $\rho = M/V$  C1
- $V = [\pi(0.38 \times 10^{-3})^2 \times 25.0 \times 10^{-2}]/4 (= 2.835 \times 10^{-8} \text{ m}^3)$  C1
- $\rho = (0.225 \times 10^{-3})/2.835 \times 10^{-8}$   
 $= 7940 \text{ (kg m}^{-3}\text{)}$  A1
- $\Delta\rho/\rho = 2(0.01/0.38) + (0.1/25.0) + (0.001/0.225) [= 0.061]$   
or  
 $\%\rho = 5.3\% + 0.40\% + 0.44\% (= 6.1\%)$  C1
- $\Delta\rho = 0.061 \times 7940 = 480 \text{ (kg m}^{-3}\text{)}$
- density =  $(7.9 \pm 0.5) \times 10^3 \text{ kg m}^{-3}$  or  $(7900 \pm 500) \text{ kg m}^{-3}$  A1 [5]
- 2 (a) (i) horizontal component  $(= 12 \cos 50^\circ) = 7.7 \text{ m s}^{-1}$  A1 [1]
- (ii) vertical component  $(= 12 \sin 50^\circ \text{ or } 7.7 \tan 50^\circ) = 9.2 \text{ m s}^{-1}$  A1 [1]
- (b)  $v^2 = u^2 + 2as$  and  $v = 0$  or  $mgh = \frac{1}{2}mv^2$  or  $s = v^2 \sin^2 \theta / 2g$  C1
- $9.2^2 = 2 \times 9.81 \times h$  hence  $h = 4.3 \text{ (4.31) m}$  A1 [2]
- alternative methods using time to maximum height of 0.94 s:
- $s = ut + \frac{1}{2}at^2$  and  $t = 0.94 \text{ (s)}$  (C1)  
 $s = 9.2 \times 0.94 - \frac{1}{2} \times 9.81 \times 0.94^2$  hence  $s = 4.3 \text{ m}$  (A1)
- or  
 $s = vt - \frac{1}{2}at^2$  and  $t = 0.94 \text{ (s)}$  (C1)  
 $s = \frac{1}{2} \times 9.81 \times 0.94^2$  hence  $s = 4.3 \text{ m}$  (A1)
- or  
 $s = \frac{1}{2}(u + v)t$  and  $t = 0.94 \text{ (s)}$  (C1)  
 $s = \frac{1}{2} \times 9.2 \times 0.94$  hence  $s = 4.3 \text{ m}$  (A1)
- (c)  $t (= 9.2/9.81) = 0.94 \text{ (0.938) s}$  C1
- horizontal distance =  $0.938 \times 7.7 (= 7.23 \text{ m})$  C1
- displacement =  $[4.3^2 + 7.23^2]^{1/2}$  C1
- $= 8.4 \text{ m}$  A1 [4]

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2016	9702	21

- 3 (a) (i) force ( $= mg = 0.15 \times 9.81$ ) = 1.5 (1.47) N A1 [1]
- (ii) resultant force (on ball) is zero so normal contact force = weight  
*or*  
the forces are in opposite directions so normal contact force = weight  
*or*  
normal contact force up = weight down A1 [1]
- (b) (i) (resultant) force proportional/equal to rate of change of momentum B1 [1]
- (ii) change in momentum =  $0.15 \times (6.2 + 2.5)$  (= 1.305 N s) C1
- magnitude of force =  $1.305/0.12$   
= 11 (10.9) N A1
- or*
- (average) acceleration =  $(6.2 + 2.5) / 0.12$  (=  $72.5 \text{ ms}^{-2}$ ) (C1)
- magnitude of force =  $0.15 \times 72.5$   
= 11 (10.9) N (A1)
- (direction of force is) upwards/up B1 [3]
- (iii) there is a change/gain in momentum of the floor M1
- this is equal (and opposite) to the change/loss in momentum of the ball so momentum is conserved A1 [2]
- or*
- change of (total) momentum of ball and floor is zero (M1)  
so momentum is conserved (A1)
- or*
- (total) momentum of ball and floor before is equal to the (total) momentum of ball and floor after (M1)  
so momentum is conserved (A1)

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2016	9702	21

- 4 (a) the energy (stored) in a body due to its extension/compression/deformation/  
change in shape/size B1 [1]
- (b) (i) two values of  $F/x$  are calculated which are the same  
e.g.  $10.4/40 = 0.26$  and  $6.5/25 = 0.26$  B1
- or
- ratio of two forces and the ratio of the corresponding two extensions are  
calculated which are the same  
e.g.  $5.2/10.4 = 0.5$  and  $20/40 = 0.5$  (B1)
- or
- gradient of graph line calculated and coordinates of one point on the  
line used with straight line equation  $y = mx + c$  to show  $c = 0$  (B1)
- (so) force is proportional to extension (and so Hooke's law obeyed) B1 [2]
- (b) (ii) 1.  $k = F/x$  or  $k = \text{gradient}$  C1
- gradient or values from a single point used e.g.  $k = 10.4/(40 \times 10^{-2})$
- $k = 26 \text{ N m}^{-1}$  A1 [2]
2. work done = area under graph  
or  $\frac{1}{2}Fx$  or  $\frac{1}{2}(F_2 + F_1)(x_2 - x_1)$   
or  $\frac{1}{2}kx^2$  or  $\frac{1}{2}k(x_2^2 - x_1^2)$  C1
- =  $\frac{1}{2} \times 10.4 \times 0.4 - \frac{1}{2} \times 5.2 \times 0.2$  C1  
or  $\frac{1}{2} \times (5.2 + 10.4) \times 20 \times 10^{-2}$   
or  $\frac{1}{2} \times 26 \times (0.4^2 - 0.2^2)$
- = 1.6 J A1 [3]
- (c) remove the force and the spring goes back to its original length B1 [1]
- 5 (a)  $T = 4 \text{ (ms)}$  or  $4 \times 10^{-3} \text{ (s)}$  C1
- $f = 1/T = 1/0.004$
- = 250 Hz A1 [2]
- (b) intensity  $\propto (\text{amplitude})^2$  and amplitude = 2.8 (2.83) (cm) B1
- curve with same period and with amplitude 2.8 cm B1
- curve shifted 1.0 ms to left or to right of wave X B1 [3]

Page 5	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2016	9702	21

- (c) (i) gradient =  $(4.5 - 2.4) \times 10^{-3} / (3.25 - 1.75) [= 1.4 \times 10^{-3}]$  B1
- wavelength =  $0.45 \times 10^{-3} \times 1.4 \times 10^{-3}$  C1
- =  $6.30 \times 10^{-7} \text{ (m)}$  C1
- = 630 nm A1 [4]
- (ii) (gradient is equal to  $\lambda/a$  therefore) gradient of line is reduced B1
- value of  $x$  will be reduced for all values of  $D$   
or new line is completely below old line  
or intercept is less B1 [2]
- 6 (a) (coulomb is) ampere second B1 [1]
- (b) (total) charge or  $Q = nAle$  M1
- $I = Q/t$  and  $l/t = v$  M1
- $I = nAle/t = nAve$  therefore  $v = I/nAe$  A1 [3]
- (c) (i) ratio =  $(I/nA_Ye)/(I/nA_Ze)$  C1
- =  $A_Z/A_Y$  or  $4A/A$  or  $\pi d^2/(\pi d^2/4)$  C1
- = 4 A1 [3]
- (ii)  $R = \rho l/A$  or  $R = 4\rho l/\pi d^2$  B1
- $R_Y = \rho l/A$  and  $R_Z = \rho(2l)/4A$  so  $R_Y/R_Z = 2$   
or  
 $R_Y = 4\rho l/\pi d^2$  and  $R_Z = 4\rho(2l)/\pi 4d^2$  or  $2\rho l/\pi d^2$  so  $R_Y/R_Z = 2$  A1 [2]
- (iii)  $V = 12R_Y/(R_Y + R_Z)$  or  $I = 12/(R_Y + R_Z)$  and  $V = IR_Y$  C1
- $V = 12 \times 2/3$
- = 8(.0)V A1 [2]
- (iv) ratio =  $I^2R_Y/I^2R_Z$  or  $(V_Y^2/R_Y)/(V_Z^2/R_Z)$  or  $(V_YI)/(V_ZI)$
- = 2 A1 [1]

Page 6	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2016	9702	21

- 7 (a) hadron: neutron/proton  
*and*  
 lepton: electron/(electron) neutrino B1 [1]  
*(allow other correct particles)*
- (b) (i) proton: up up down or uud B1 [1]  
 (ii) neutron: up down down or udd B1 [1]
- (c) (i) neutron  $\rightarrow$  proton + electron + (electron) antineutrino B1 [1]  
 (ii) up down down (quarks) change to up up down (quarks)  
*or*  
 down (quark) changes to up (quark) B1 [1]